

University of Dundee

How Plants Grow Up

McKim, Sarah

Published in:
Journal of Integrative Plant Biology

DOI:
[10.1111/jipb.12786](https://doi.org/10.1111/jipb.12786)

Publication date:
2019

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):
McKim, S. (2019). How Plants Grow Up. *Journal of Integrative Plant Biology* , 61(3), 257-277.
<https://doi.org/10.1111/jipb.12786>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Invited Expert Review

How Plants Grow Up

Running Title: How Plants Grow Up

Sarah M McKim

Division of Plant Sciences, University of Dundee at The James Hutton

Institute, Invergowrie, Dundee DD2 5DA, UK

***Correspondence:** Sarah M McKim (s.mckim@dundee.ac.uk)

Edited by: Thorsten Schnurbusch, Leibniz Institute of Plant Genetics and
Crop Plant Research (IPK), Germany

This is the peer reviewed version of the following article: McKim, S. "How Plants Grow Up", *Journal of Integrative Plant Biology* (2019) which has been published in final form at <https://doi.org/10.1111/jipb.12786>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

This article is protected by copyright. All rights reserved.

Received: October 15, 2018; Accepted: January 21, 2019

ABSTRACT

A plant's lateral structures, such as leaves, branches and flowers, literally hinge on the shoot axis, making its integrity and growth fundamental to plant form. In all plants, subapical proliferation within the shoot tip displaces cells downwards to extrude the cylindrical stem. Following the transition to flowering, many plants show increased axial elongation with associated increases in subapical proliferation and expansion. However, the stems of grasses, called culms, also elongate due to detached intercalary meristems, which displace cells upwards to elevate the grain-bearing inflorescence. Variation in culm length within species is especially relevant to cereal crops, best exemplified by the high-yielding semi-dwarfed cereals in the Green Revolution. While pathways controlling subapical and intercalary stem growth have been understudied, renewed interest in axial development reveals that control of cell division planes, boundary formation and temporal dynamics of differentiation, are likely critical mechanisms coordinating axial growth and development in plants.

INTRODUCTION

EVOLUTIONARY ORIGINS OF THE LAND PLANT STEM

The central cylindrical axis of land plants is a deeply conserved and ancient feature, shared with their bryophyte-like ancestors and predating roots and leaves. In early land plants, stems were diminutive stalks for the terminal sporangium acting to aid spore dispersal (Cronk 2009; Ligrone et al. 2012). The advent of lignified vasculature allowed more extensive vertical growth by providing mechanical support and conductive channels for solute transport over long distances. Accessing the higher canopy helped sporophytes better evade soil-borne pathogens and diversify their reproductive strategies. Axial growth also increased photosynthetic capacity in the aerial space which, especially following the evolution of 3D growth, contributed to sporophyte dominance.

STEM MORPHOGENESIS IN VASCULAR PLANTS

Lateral 3D growth from the shoot tip generates leaves and branches which attach to the vertical axis at nodes, segmenting the stem into alternating nodes of lateral organ attachment and intervening internodes. The shoot is thus organised into iterative units called phytomers which comprise a node, its lateral organ, subtending internode and axillary bud (Galinat 1959; Bossinger 2009). All these structures derive from self-renewing indeterminate cells and proliferation at the shoot tip. Thus, unlike the basal intercalary proliferation which lifts the moss sporophyte, the vascular plant stem elongates due to downward displacement of daughter cells to form a “self-extruding cylinder” (Cronk 2009; Harrison and Morris 2018).

Cells in the shoot tip make up the shoot apical meristem which is organised into distinct, functional zones (Figure 1A). The central apical zone consists of a small number of stem cells which is surrounded by the more mitotically-active peripheral zone whose cells are recruited to form lateral organs. The underlying stem grows from two subapical regions: the ‘rib’ meristem basal to the central zone, whose transversely dividing cells form the stem’s central core (pith); and from subapical peripheral cells contributing the stem epidermis and cortex (Vaughan 1955; Sachs et al. 1959b; Sachs 1965; Kwiatkowska 2008; Bencivenga et al. 2016; Greb and Lohmann 2016). Peripheral and central subapical rib regions, including the rib meristem, together constitute the rib zone (Figure 1A; Gaillochet et al. 2015; Bencivenga et al. 2016; Janocha and Lohmann 2018). However, all of these cells have also been called the rib meristem (Perales and Reddy 2012), while sometimes the rib zone is shown separate from the rib meristem (Fletcher 2002). Furthermore, defining the basal borders of the rib zone is challenging since

the axial transition from dividing to differentiating cells lacks an obvious boundary while the length of this region itself varies by species and growth stage (Sachs 1965; Fisher and French 1976). The distinctive phytomeric production in grasses compared to dicots also impacts this subapical zonation (Figure 1B). In dicots, the lateral primordium (leaf or flower bud) bulges from a triangular top flank off the peripheral zone, while in grasses, the leaf primordium remains connected to a basal ring of proliferating founder cells circumnavigating the shoot tip, called the 'disc of insertion' (Sharman 1942). While the leaf blade develops from the lateral primordium growing away from the apex, the upper disc of insertion develops into the leaf sheath while the lower disc of insertion forms the node, internode and axillary bud (Sharman 1942; Johri and Coe 1983; McDaniel and Poethig 1988; Scanlon and Freeling 1997). In this way, the grass stem represents an iterative stack of clonally-related phytomeric units.

SECONDARY AXIAL ELONGATION

Sachs (1965) recognised two phases of stem development: primary stem morphogenesis, which establishes radial patterning and the node-internode arrangement, and secondary elongation, that occurs in some species depending on growth stage and environment. Secondary elongation is distinct from the secondary thickening of dicot stems due to vascular cambium proliferation (reviewed in Sanchez et al. 2012). A classic example of secondary elongation occurs following flowering in the crucifers, Pooideae grasses (including the cereal crops) and other species which undergo extensive internode elongation that shifts their growth habit from compressed to caulescent (often referred to as 'bolting'; Figure 1C, D; Sachs 1965; Kellogg 2015), elevating the inflorescence into the upper canopy to promote pollen exchange, seed dispersal and evade pathogens. Secondary elongation can influence both the length of extant internodes, as well as those generated after the reproductive transition. For instance, *Arabidopsis* stems develop long internodes between nodes projecting axillary branches subtended by leaves (cauline paraclades), and then elongated internodes separating solitary flowers, each on their own pedicels, which continually emerge on an exposed inflorescence. In Pooideae grasses, the vegetative internodes elongate and are supported by the leaf sheaths until the final internode extends to push the flowering head out of the final leaf sheath, exposing the developed inflorescence, usually at anthesis. Emergence, called 'heading', is critical to harvest the grain-laden inflorescence and must be tightly regulated as overly tall culms are prone to lodge or fall over, thereby leading to lost yield, making internode elongation a key trait for yield. Semi-dwarf varieties lodge less, which contributed to the vast yield improvements during the Green Revolution.

Increased mitotic activity and/or size of cells in the rib zone occurs during secondary axial elongation in both dicots and monocots (Figure 1C, D; Sachs et al. 1959b; Fahn et al. 1963; Jacqumard et al. 2003; Kwiatkowska 2008). However, while not a uniform feature of monocots (Sachs 1965; Fisher and French 1976), individual grass internodes elongate due to intense cell division within an 'intercalary meristem' just above each node: here, transverse divisions displace cells upwards through successive zones of expansion and maturation, lengthening each internode from the bottom up (Figure 1D; Schmalfluss 1930; Evans 1965; Kaufman et al. 1965; Fisher 1970; Fisher and French 1976; Bleecker et al. 1986; Cho and Kende 1997; Martin et al. 2016). Intercalary growth gives rise to most of the cells in the culm, which is typically formed of internal cortex and inner stem tissues of scattered vascular bundles and supporting sclerenchyma, contained within an epidermis of axially arranged cell files, including files of stomata, cork and silica cells (Harting 1845; Grisebach 1943; Kaufman et al. 1965). Intercalary meristems are only found in monocots with sheathing leaf bases (Fisher and French 1978) that start elongating before their corresponding internode in an acropetal pattern (Kirby et al. 1994), to provide mechanical support to fragile, meristematic and actively elongating internode tissues both within their own and more apical internodes (Niklas 1990). While the intercalary meristem is still active, internodes in some grass species develop a central cavity or lacuna early during elongation due to comparatively slow pith growth, leading to the characteristic hollow straw of wheat, rice and barley. Internodes in other species remain solid as in maize and sugarcane, or a combination of solid and hollow as in bamboo (Kellogg 2015). However, nodes always remain solid and become thickened with a complex vascular plexus of anastomosed vascular tissue from the stem, leaves and buds (Sharman and Hitch 1967; Kellogg 2015). Thus, an important distinction between bolting of *Arabidopsis* and internode growth of grasses following flowering is the developmental origin of the internode. In *Arabidopsis*, cells in the bolting internodes are fueled exclusively cells from the rib meristem after the reproductive transition (Vaughan 1955; Jacqumard et al. 2003) while dividing cells of the grass intercalary meristems either are or quickly become detached from subapical proliferation by differentiating internode tissue. Furthermore, the intercalary meristem, whose activity declines over time (Kaufman et al. 1965), is determinate and generates a single internode. In this way, the intercalary meristem is similar to the leaf plate meristem, a set of parallel cell layers at the base of leaves whose anticlinal divisions contribute to leaf outgrowth; in contrast, the rib meristem usually has an indeterminate nature and continuously divides.

As with other organs, final internode length reflects cell number and the extent of longitudinal cell elongation. GA promotes both intercalary and rib meristem-driven secondary axial elongation through increasing cell division

and cell elongation (Sachs et al. 1960; Sachs 1965; Kende et al. 1998). Interfering with this process leads to varying degrees of dwarfism, which in the cultivated cereals, higher yields due to decreased partitioning into vegetative biomass and reduced lodging. For instance, the semi-dwarf rice and wheat of the Green Revolution are altered in GA biosynthesis or perception, respectively (Webb et al. 1998; Peng et al. 1999; Hedden 2003). The *Reduced height* wheat semi-dwarfs show reduced responsiveness to GA (Gale and Youssefian 1985) due to truncated, constitutively-active DELLA transcription factors (Peng et al. 1999), negative regulators of GA signalling, normally suppressed by GA (Peng et al. 1997). In rice, the *semidwarf-1* locus encodes a GA 20-oxidase (GA20ox) enzyme critical for GA biosynthesis, which when defective leads to semi-dwarfism (Monna et al. 2002; Sasaki et al. 2002; Spielmeyer et al. 2002). Other semi-dwarfing alleles derive from defects in brassinosteroid (BR) growth hormone biosynthesis or perception/signalling (Chono 2003; Dockter et al. 2014; Hirano et al. 2017). For instance, in barley, the BR insensitive mutant *uzu* has a single nucleotide substitution in the gene encoding the BR receptor, *HvBRI1* (Chono et al. 2003), leading to a temperature-sensitive allele (Dockter et al. 2014). Dockter et al. (2014) also described mutant alleles of three BR biosynthetic genes – *HvBRD* (*BR 6-OXIDASE*), *HvCPD* (*BRASSINOSTEROID C-23 HYDROXYLASE*) and *HvDIM* (*DIMINUTO*), which each encode enzymes of the BR pathway, and display similar phenotypes, including erect and small leaves as well as reduced stature. These features of BR semi-dwarfs are usually attributed to reduced cell elongation as opposed to cell proliferation (Szekeres et al. 1996; (Choe et al. 1999; Sun et al. 2010); however, loss of a function of BRI1 in rice leads to shorter internodes associated with improper formation of the IM (Yamamuro et al. 2000). Many roles of BR may reflect molecular and feedback interactions with GA metabolism and signalling to control growth (Bai et al. 2012; Gallego-Bartolomé et al. 2012; Li et al. 2012; Tong et al. 2014) but recent analyses in maize suggest that the extent of interaction between the two hormones depends on the developmental context (Best et al. 2016).

Several reviews comprehensively discuss semi-dwarfing alleles in cereals (Hedden 2003; Dockter and Hansson 2015; Braumann et al. 2018). Unsurprisingly, many other genes whose alleles influence stem elongation are associated with changes in hormone metabolism and/or signalling. An important task is to understand how these alleles participate in the regulatory pathways influencing how, where and when these growth hormones function during primary stem development and secondary stem elongation across plants. In their recent review, Serrano-Mislata and Sablowski (2018) expertly describe how new approaches to study stem organogenesis revealed the importance of oriented growth in the rib zone as well as the role of cell wall mechanics during cell expansion in stem elongation and integrity. Here, I attempt to summarise what we know about the genetic mechanisms controlling the balance between proliferation and differentiation during primary stem development with an emphasis on grass models and Arabidopsis, and discuss the relevance of these mechanisms to secondary intercalary activity,

as well as control of secondary axial elongation. I conclude with outstanding questions deserving attention.

REGULATION OF SUBAPICAL PROLIFERATION

The leaf and subtending internode derive from the same cell population (Jegla and Sussex 1989; Poethig and Szymkowiak 1995). Mutants defective in apex maintenance often exhibit aberrant stem morphology, uneven internode lengths and disturbed phyllotaxis, suggesting a coordination of lateral organogenesis with stem morphogenesis at the shoot tip. Below, I describe several central pathways associated with meristem function and consider their influence on stem development.

Iterative and modular plant growth depends on maintaining a stable meristematic cell population despite continual depletion of daughter cells into new phytomers. The CLAVATA (CLV) – WUSCHEL (WUS) signalling loop is key to this balance, and although best understood in *Arabidopsis*, the pathway's principles appear broadly conserved across plants (reviewed in Somssich et al. 2016; Figure. 2A). The *WUS* gene encodes a homeodomain whose loss of function leads to arrested growth due to meristem consumption by lateral organogenesis (Laux et al. 1996). This reflects a role of *WUS*, expressed in the rib meristem (a pattern also described as marking an organising centre), to non-cell autonomously promote proliferation and stem cell identity in overlying central cells and restricts expression of differentiation-associated genes to the meristem periphery (Yadav et al. 2014; Gruel et al. 2016). *WUS* also stimulates the overlying cells to produce CLV3, a mobile peptide ligand, which when perceived by Leucine Rich Receptor (LRR)-receptor complexes surrounding and within the rib meristem, signals repression of *WUS* transcription, thereby completing a negative feedback loop (Schoof et al. 2000; Brand et al. 2002; Muller et al. 2006; DeYoung and Clark 2008; Yadav et al. 2009, 2013; Yadav and Reddy 2011; Snipes et al. 2018). Disruption of CLAVATA function uncouples proliferation from organogenesis, causing expanded/ ectopic *WUS* expression domains and ever-increasing stem cell proliferation. Associated with altered cell division planes, ectopic *WUS* leads to profound changes in meristem shape and degrees of fasciation (Leyser and Furner 1992; Medford et al. 1992; Clark et al. 1993, 1996; Kayes and Clark 1998; Suzaki et al. 2004; Bommert et al. 2005; Suzaki et al. 2006; Xu et al. 2015; Mandel et al. 2016). Fasciation occurs as the apical meristem progressively widens, flattens and divides into connected meristem ridges, a change in shape and function propagated downwards to form a bundled, fused stem (Cronk 2009). Mutants defective in cell cycle also show fasciation, such as in the maize *tangled* severe dwarf (Smith et al. 1996; Kaya et al. 2001; Suzaki 2004) while altered frequency and orientation of meristematic

cell divisions are also associated with altered *WUS* expression in *Arabidopsis* (Kaya et al. 2001; Suzaki et al. 2004; Muñoz-Nortes et al. 2014), emphasising the importance of controlling the pattern of *WUS* expression for proper axial growth. The normal, single central subapical localisation of *WUS* expression is established in the embryo and stably maintained through cytokinin signalling (Leibfried et al. 2005; Gordon and Chickarmane 2009; Chickarmane et al. 2012; Adibi et al. 2016; Meng et al. 2017; Snipes et al. 2018). *WUS* induction of *HECATE* transcription factor genes, whose activity reduces auxin signalling, is critical for meristem integrity (Schuster et al. 2015; Gaillochet et al. 2018), consistent with reports that both auxin treatment and inhibition of auxin transport cause fasciation (Gorter 1965; Fambrini et al. 2006). Taken together, regulated cell division orientation associated with a centrally defined *WUS* expression domain and corresponding hormonal balance, are essential for cylindrical axis integrity and corresponding stem growth.

Fasciated mutants also show aberrant phyllotaxy, uneven internode growth and dwarfing, suggesting that meristem integrity is also key to coordinate stem growth with lateral organ emergence (Smith and Hake 2003; Running et al. 2004; Khan et al. 2012; Pautler et al. 2015). However, these phenotypes are not always present together and can be restricted to a particular growth stage or meristem type, likely reflecting both the context-dependent nature of individual *WUS*-CLV signalling pathways. Involving multiple *CLAVATA3/ENDOSPERM SURROUNDING REGION* (*CLE*) peptides and their CLV-like receptors and co-factors, as well as multiple *WUS*-like *HOMEBOX* (*WOX*s), these pathways control proliferation in specific contexts, although many intervening steps and exact protein/peptide partners are not completely understood, especially in grasses (Pautler et al. 2013; Somssich et al. 2016; Bommert and Whipple 2018).

For instance, we don't know whether apical CLV3-like signals control subapical proliferation exclusively through downward displacement of peripheral cells or whether cells subapical or peripheral to the rib meristem themselves emit additional signaling *CLE*s. In *Arabidopsis*, several *CLE*s are expressed beneath the *WUS* expression domain, yet none show a loss-of-function phenotype, perhaps due to overlapping expression and functional redundancy (Jun et al. 2010; Gregory et al. 2018). More is known about mechanisms delineating the *WUS* basal boundary in grasses. Rather than sole expression in a central organising or rib meristem, maize vegetative apices transiently express *ZmWUS1* within the lower disc of insertion containing the presumptive node, internode and axillary bud cells (Figure 2B; Nardmann and Werr 2006). Je et al (2016) showed that perception of a leaf primordium-derived *CLE* peptide *ZmFON2-LIKE CLE PEPTIDE1* (*ZmFCP1*) by the *FASCIATED EAR3* (*FEA3*) CLV-like receptor, which is expressed below and inside the *ZmWUS1* region, prevents basal expansion of *ZmWUS1*

expression in the transitional inflorescence meristem (Je et al. 2016); however, it is unclear if this mechanism also limits later *ZmWUS1* expression in the developing phytomer. Arabidopsis *fea3* mutants are short and fasciated, while *clv1* mutants in Arabidopsis (*clv1*, Leyser and Furrer 1992), maize (*thick dwarf tassel1*, *td1*, Bommert, 2005), rice (*floral organ number1*, Moon et al. 2006) and cucumber (*Cscclv1*, Xu et al. 2018) are dwarf; however, neither Arabidopsis *clv3*, *clv2* or maize *fea3* mutants display altered plant height (Clark et al. 1995; Je et al. 2016). Partial expression overlap in the meristem flank and the lateral organ domains *ZmWUS2*, the other *WUS*-like gene in maize, and *TD1*, suggests a possible control of TD1 over the *ZmWUS2* domain, potentially impacting the entire phytomer (Nardmann and Werr 2006). Impaired expression or function of other LRR-like kinase receptors such as BARELY ANY MERISTEMS (BAMs) and ERECTA(ER)/ERL, can enhance *clv* fasciation and dwarfism in Arabidopsis (Yokoyama et al. 1998; DeYoung and Clark 2008), although the function of grass orthologues are unknown. Fine-tuning of receptor complexes via their spatiotemporal levels and interacting partners may integrate diverse peptide signals (Janocha and Lohmann 2018), which possibly explains the severe phenotypes of certain LRR mutants.

In the grasses, a subset of peripheral phytomer founder cells specify a shared marginal domain in the leaf and internode, which when absent leads to narrow leaves, lacking margins and subtending internodes shorter on their side adjacent to the leaf sheath margin (Scanlon et al. 1996). Cloning the *narrow sheath1/2* (*ns1/2*), *narrow leaf2/3* and *narrow leafed dwarf1* mutants in maize, rice and barley, respectively, identified the *WOX3*-like genes as essential to recruit marginal domain founder cells, orthologous to the *PRESSED FLOWER* (*PRS*) gene of Arabidopsis necessary for marginal development in the flower (Figure. 2B; Matsumoto and Okada 2001; Nardmann 2004; Cho et al. 2013; Yoshikawa et al. 2016). However, while neither rice *narrow leaf2* or 3 mutants show differences in plant height, overexpression of *OsWOX3A* in rice led to extreme dwarfism and marginal expansion (Ishiwata et al. 2013), recently associated with downregulation of GA biosynthesis, likely through directly interaction with a GA biosynthetic gene promoter (Cho et al. 2016). Thus, marginal identity must be carefully regulated to recruit sufficient founder cells but balanced with differentiation capacity, which may be related to hormonal homeostasis amongst meristem, founder and differentiating tissues.

Potential partners linking meristem maintenance and stem elongation, especially in response to environmental cues, are heterotrimeric G-proteins which interact with CLV signalling (reviewed in Urano et al. 2016). A major dwarfing locus in rice, *DENSE PANICLE1* (*DEP1*) and barley (*HvDEP1*) encodes a G γ subunit (Huang et al. 2009; Wendt et al. 2016; Xu et al. 2016) while defective alleles in genes encoding the G α subunit in rice ('*daikoku*' *dwarf1*, Fujisawa et al. 1999; Ashikari et al. 1999), barley (*brachytic1*, Ito et al. 2017; Braumann

et al. 2018b), maize (*compact plant2*, *ct2*, Bommert et al. 2005) and Arabidopsis (*gpa1*, Ullah, 2001) confer compact shoots due to reduced axial proliferation. In rice, *DWARF1* interacts with GA and BR signalling to control height (Ashikari et al. 1999; Oki et al. 2009). In Arabidopsis, the G-protein β -subunit1 (AGB1) works synergistically with a CLV receptor in meristem maintenance (Ishida et al. 2016), while maize *CT2*, expressed throughout the apex and emerging leaves, signals downstream of FEA2, a CLV2 receptor, as well as independently to control stem cell proliferation (Bommert et al. 2013), highlighting the involvement of heterotrimeric G proteins within the CLV-WUS pathway.

Does a CLV-WUS pathway regulate maintenance of intercalary meristems during secondary stem elongation? Intercalary meristems maintain their integrity prior to their activation and continue dividing for an extended period during internode growth (Kaufman et al. 1965; Bleecker et al. 1986). It is long known that dwarf grass plants usually have fewer cells (Harting 1843), suggesting inadequate intercalary proliferation, as well as the existence of mechanisms ensuring either sufficient founder cell recruitment and/or proliferative capacity in the intercalary meristem. Since expression of *WOX3/PRS* in the nascent internode appears essential for internode growth on the marginal side, these genes could also regulate internode cell recruitment and/ or marginal intercalary proliferation. Another candidate is the *DWARF TILLER (DWT)* locus in rice, encoding an *OsWOX* necessary for cell proliferation and elongation and GA signalling; however, loss of function leads to dwarfed tillers only, and *DWT* is not expressed in the internode, suggesting a non-cell autonomous regulation of intercalary growth in tillers (Wang et al. 2014). Other candidates are the *WOX3* clade sister to the narrow leaf *PRS/WOX3*'s which arose due to an ancient duplication in grasses (Nardmann et al. 2007). Unlike marginal domain-specific expression domain of *WOX3/PRS* (Nardmann et al. 2004; Shimizu et al. 2009), sister *WOX3*s are expressed in a complete ring around the initiating phytomer which is then restricted to the phytomer base (Nardmann et al. 2007), and in rice, are strongly expressed later on in the intercalary region and nodal vasculature of young internodes (Figure 2C; Angeles-Shim et al. 2012), suggesting subfunctionalisation of the sister *WOX3* clade for the distinctive structures of grasses. However, to date, only the rice sister *WOX3* has been characterised, and reported to control trichome formation (Angeles-Shim et al. 2012; Zhang et al. 2012; Yoshikawa et al. 2016) so the relevance of this expression pattern is unclear.

Overlying the CLV-WUS pathway are the Class I KNOTTED-like homeoboxes (KNOX-I) whose promotion of meristem activity is conserved across vascular plants (Harrison et al. 2005). *KNOX-I* expression (Figure 2C), often used as a marker for meristematic identity, promotes proliferation in meristems by

upregulating cytokinin biosynthesis (Jasinski et al. 2005; Yanai et al. 2005), downregulating GA biosynthesis (Hay et al. 2002) and antagonising expression of differentiation-associated gene (Byrne et al. 2002). Dramatic downregulation of *KNOX-I* marks presumptive primordium cells and is required for lateral organ identity and differentiation (Scanlon 2003). Loss of the marginal domain in maize *ns* mutants is associated with a failure to downregulate *KNOX-I* (Scanlon et al. 1996), suggesting that founder cell recruitment into the emerging internode also depends on *KNOX-I* downregulation. This role is consistent with both the dwarfism caused by the dominant overexpression mutations in maize for *Knotted1* (*Kn1*; Freeling and Hake 1985; Vollbrecht et al. 1991), *Gnarley/KNOX4* (*Gn1*, Foster et al. 1999) and *Rough sheath1* (*Rs1*, Becraft and Freeling 1994) and the direct upregulation of a GA catabolism gene (*ga2ox1*) by *KN1* (Bolduc and Hake 2009). However, loss of function *KNOX-I* alleles also lead to dwarfism in rice (*osh15*, Sato et al. 1999); *osh1*, Tsuda et al. 2011), maize (*rs1*, Schneeberger et al. 1995) and Arabidopsis (*brevipedicellus*, *bp*, Venglat et al. 2002), potentially due to loss of subapical meristematic proliferation. *KNOX-I* show persistent expression in regions forming the stem (Reiser et al. 2000), such as *BP* in the rib meristem and growing stem vasculature of Arabidopsis (Lincoln et al. 1994; Smith and Hake 2003), *Kn1* and *Rs1* in the rib zone and lower disc of insertion of maize, overlapping with *ZmWUS1* (Jackson et al. 1994; Schneeberger et al. 1995; Nardmann and Werr 2006) and similar to *OSH15* localising to the base of growing leaves (Sato et al. 1999), while *Kn1* is expressed throughout intercalary meristems (Tsuda et al. 2017). Coupled with their mutant phenotypes, *KNOX-1* expression patterns suggest important roles in both subapical and intercalary proliferation.

REGULATION OF BOUNDARY IDENTITY AND VASCULAR DIFFERENTIATION

The *osh15* and *bp* mutants also show aberrant, ectopic and/or excessive internode lignification, which may mechanically hamper elongation (Sato et al. 1999; Smith and Hake 2003). While both *BP* and *OSH15* directly inhibit expression of lignin biosynthetic genes (Mele et al. 2003; Yoon et al. 2017), several studies also suggest that antagonistic interactions between *KNOX-I*'s themselves, and interaction with *BEL-LIKE HOMEODOMAIN* (*BLH*) genes (*BLHs*), influence vascular development and boundary positioning. In Arabidopsis, *BP* physically interacts with the *PENNYWISE/REPLUMLESS* (*PNY/RPL*), a *BLH* whose loss of function resembles *bp*, while double *bp pny* mutants are severely dwarfed with a thick continuous vascular ring in their stems (Smith and Hake 2003). Similar phenotypes occur from overexpression of boundary marking genes, including *KNOX-I* genes *KNAT2/6*, the *BLH* gene *ARABIDOPSIS THALIANA HOMEODOMAIN1* (*ATH1*) and the *BLADE-ON-PETIOLE* genes (Smith and Hake 2003; Norberg et al. 2005; Cole et al. 2006;

Ragni et al. 2008; Gómez-Mena and Sablowski 2008; Khan et al. 2012). *ATH1* is particularly interesting since impaired *ATH1* leads to longer internodes and an expansion of the rib zone based on *BP* expression, suggesting that *ATH1* represses rib zone proliferation (Gomez-Mena and Sablowski 2008). Ectopic boundary gene expression causes excess lignification, altered phyllotaxy, dwarfism and uneven internode length in *bp* and *pny* since loss of function boundary gene alleles rescues these phenotypes (Ragni et al. 2008; Khan et al. 2012; Bencivenga et al. 2016). Thus, *BP* and *PNY* likely interact to exclude expression of these boundary genes from the developing internode, limiting their expression to the boundary (Figure 3A). These roles may originate very early during stem morphogenesis (Figure 3A): in their insightful study, Bencivenga et al (2016) showed that loss of *PNY* lead to medial expansion of boundary gene expression from the peripheral to central rib zone region, which decreased transverse cell divisions and correspondingly increased more peripheral-like radial divisions in the central rib zone. Accordingly, *PNY* may prevent peripheral rib zone encroachment into the central rib region through restricting boundary gene expression to the periphery, similar to *PNY*'s role in defining the boundary between the central and peripheral apical meristem zones (Ung et al. 2011). Since stem vasculature in *Arabidopsis* develops at the interface between the central and peripheral rib zone regions, Bencivenga et al (2016) speculate that altered cell orientations could influence this positioning, leading to the increased number of vascular strands in *pny*. Altered peripheral and central positioning may also contribute to the thick continuous vascular ring typical of *bp pny* stems.

Regulation of internode elongation by specification and/or proliferation of stem vasculature appears conserved in maize, despite its different proliferation and vascularisation pattern compared to dicots (Figure 3B). Exciting recent work by Tsuda et al (2017) showed that loss of function in *BLH12* and *BLH14* (maize *BLHs* of the *PNY* clade), led to truncated internode elongation, due to earlier arrest of intercalary meristem activity as well as premature vein anastomoses. Since *BLH12*, *BLH14* and *KN1* proteins interact and are all detected in the intercalary meristem, suggesting that *BLH12/14* and *KN1* may work together to maintain intercalary meristems and prevent their precocious differentiation (Tsuda et al. 2017). *BLH12* and *BLH14* as well as *KN1* also accumulate in the provasculature (Tsuda et al. 2017; Jackson et al. 1995), where they slow the timing of the progressive lignification program in internodes. Downregulation of *BR* via by *OSH1* is also important for correct boundary formation and positioning (Tsuda et al. 2014). Since *BR* also promotes xylem differentiation (Caño-Delgado et al. 2004), it is plausible that increased/ ectopic vascularisation due to loss of *KNOX-1*, *BLH* may reflect increased *BR* leading to premature differentiation and vascularisation. Taken together, work in both dicots and monocots highlight the critical role for

KNOX-I and BLH in coordination of proliferation with vascular development and internode elongation, by controlling temporal and spatial dynamics of cell division and vascular differentiation in the internode.

The master shoot identity regulators, *CLASS III HOMEODOMAIN - LEUCINE ZIPPER (HD-ZIP III)* genes are also implicated in the control of stem growth through their effects on proliferation and vascularisation. Expression of the HD-ZIP III *ATHB8* marks preprovascular strands while its overexpression caused precocious vascular differentiation, slower stem elongation and dwarfism (Baima et al. 1995, 2001). In Arabidopsis, *ATHB8* is induced by auxin and negatively regulated by *ACAULIS5 (ACL5)*: a gene which encodes an enzyme synthesising thermospermidine (Takechi et al. 2008; Baima et al. 2014), whose loss of function *acl5* alleles lead to thicker veins and short stems (Clay and Nelson 2005). In fact, all five *HD-ZIP III* genes affect plant stature in Arabidopsis, and their higher order mutant phenotypes, which include severe dwarfing, fasciation and increased *WUS* expression, suggest they have redundant roles to ensure radial symmetry, elongation and vascular patterning in the stem (Emery et al. 2003; Prigge et al. 2005; Williams et al. 2005). In rice, these genes are also important for meristem maintenance but also for marginal domain development (Itoh et al. 2008), and can affect height as overexpression of the rice *HD-ZIP III OsHOX32* lead to semi-dwarfism, associated with changes in *YABBY1 (YAB1)* expression, a gene involved in lateral differentiation and polarity (Li et al. 2016). Less is known about the roles of *Class I HD-ZIPs*; however, *OsHOX4* gene in rice promotes GA degradation and overexpression leads to semi-dwarfism, associated with elevated expression of *YAB1* and a GA catabolism gene, *GA2ox3* (Dai et al. 2008; Zhou et al. 2015). Another *HD-ZIP I* gene, *OsHOX12*, has an opposite function and promotes stem elongation in rice by repressing the expression of *ELONGATED UPPER INTERMODE (EUI, CYP714D1)*, which encodes a GA-deactivating enzyme (Zhu et al. 2006; Gao et al. 2016).

A separate CLV-WUS pathway also influences stem morphology through effects on vasculature proliferation. Overexpression of *CLE42* and *CLE44* in Arabidopsis led to bushy, dwarf plants (Strabala et al. 2006). These peptides are now known to activate PHLOEM INTERCALLATED WITH XYLEM (PXY) receptors to restrict the expression of vascular proliferation-promoting *WOX4* and *WOX14* to the vasculature, in part through control of cell division orientation (Etchells and Turner 2010; Etchells et al. 2013). Similar to Arabidopsis, *WOX4* in rice (*OsWOX4*) is expressed in the nascent vasculature, in addition to the meristem and leaf primordia, and RNAi-mediated knock-down of *OsWOX4* was associated with stunted shoots (Ohmori et al. 2013).

It will be interesting to further unpick the interactions amongst HD-ZIPs, KNOX, BLH and CLE/WOX which control division and vascular differentiation events in elongating internodes in both monocots and dicots. Several transcriptomic datasets describing gene expression along developing grass and Arabidopsis internodes may help construct networks regulating tissue histogenesis and metabolism along this developmental gradient (Ehlting et al. 2005; Cui et al. 2012; Martin et al. 2016; Kebrom et al. 2017). Of particular interest will be comparing these networks to those underlying sporophyte elongation in non-vascular plants, which may reveal the regulatory innovations developed by vascular land plants to balance lignification with proliferation and indeterminacy to both enable axial elongation and apical indeterminacy.

TRANSITION TO SECONDARY STEM ELONGATION

Bolting in Arabidopsis occurs after the reproductive transition (Hempel and Feldman 1994), and similarly, grass culm internode elongation occurs only at specific stages of reproductive spike development across genotypes and environmental conditions (Nicholls and May 1964). Thus, internode elongation may be tuned to an apex-derived signal. In oats, internode elongation was reduced by removal of the inflorescence and lost completely when nodes were eliminated, suggestive of an apex-originated signal transported through the nodes (Koning et al. 1977). In pea and tobacco, auxin promoted GA accumulation in stems (Ross et al. 2000; Wolbang and Ross 2001), whereas in barley, inflorescence-derived auxin appears necessary for GA biosynthesis in the internode, suggesting a conserved relationship between apical auxin and GA-driven internode elongation (Wolbang et al. 2004). Furthermore, Arabidopsis, pea and other dicots, and grasses share an acropetal pattern of internode elongation with basal internodes elongating first (Kirby et al. 1994; Pouteau and Albertini 2011). As basal internodes are older, the acropetal pattern could arise if internodes acquire a particular competence for elongation based on developmental stage. Alternatively or additionally, internodes may respond to a signal which itself is acropetally distributed. GA is a promising candidate. Long known to promote both bolting and intercalary elongation when applied to internodes (Sachs 1965; Kaufman 1965; Bleecker et al. 1986), a leaf-derived acropetal GA signal appears important for internode elongation and flowering in tobacco (Eriksson et al. 2006; Dayan et al. 2012). In addition, dicot stems transport GA acropetally through the stems (Binenbaum et al. 2018). Furthermore, internodes in constitutive GA-signalling slender mutants in Arabidopsis, rice and barley concurrently elongate, a similar response observed after saturating GA treatment (Foster 1977; Ikeda et al. 2001; Aoki et al. 2002), suggesting that misregulated GA levels can uncouple the acropetal internode pattern. Interestingly, work in rice suggests differential regulation of GA level and signalling between the basal versus upper unelongated internodes: as suggested by the locus name, the enzyme

encoded by *EUI* described above appears to deactivate GA specifically in the final internode of rice (Zhu et al. 2006), while a gibberellin 2-oxidase, which also deactivates GA, encoded by the *SHORTENED BASAL INTERNODES* gene, is specifically expressed in basal internodes of rice where it prevents basal internode expansion (Liu et al. 2018). Taken together, these data suggest that regulation of GA levels, transport, perception and signalling could influence the acropetal wave of internode elongation along the shoot axis, potentially in response to a basipetal auxin signal. BR is also suggested to move acropetally in wheat and cucumber (Nishikawa et al. 1994) - although not in pea (Symons and Reid 2004) - which may interact with GA to control the elongation pattern.

Recent work by Serrano-Mislata et al (2017) using both Arabidopsis and barley showed that the master negative regulators of GA signalling DELLA proteins control apical meristem size by targeting a cell cycle inhibitor but that this function could be genetically uncoupled from its effects on secondary stem elongation. This work hints that additional DELLA targets and/or partners with a localised function to the elongating stem could be relevant for GA control of bolting growth. DELLA proteins directly interact with Class I TEOSINTE BRANCHED 1 [TB1], CYCLOIDEA [CYC], and PROLIFERATING CELL FACTOR [PCF] (TCP) proteins (Davière and Achard 2013; Davière et al. 2014) which control cell proliferation across plants (Martín-Trillo and Cubas 2010). Class I *TCP14/15* promote proliferation in internodes (Kieffer et al. 2011) and higher order class I *tcp* mutants are severely dwarfed (Davière et al. 2014). Interestingly, *tcp14 tcp15* differences originate at the apex, with axial spacing reduced between floral primordia but phyllotaxy normal, suggesting that apical meristem maintenance/size is not hugely altered (Kieffer et al. 2011). Moreover, *TCP14/15* are not expressed in inflorescence meristems and very young floral primordia but strongly expressed in young stems (Kieffer et al. 2011), so the TCPs may be local regulators mediating internode growth, similar to their role in axillary branching in dicots and monocots (Hubbard et al. 2002; Takeda et al. 2003; Aguilar-Martinez et al. 2007).

The grass internode derives from the lower disc of insertion, and remain small and developmentally delayed while the upper disc and primordium differentiate into leaves (Briggs 1978). Intercalary meristems in rice are present by the P3 stage (Kaufman 1959), suggesting that proliferating cells may enter a dormant state upon leaving the apex. The rib meristem itself is known to enter dormant states in other species: in their impressive work, Ruonala et al (2008) describe how bud dormancy in poplar is associated with arrested proliferation in the rib meristem but also callose deposition at plasmodesmata. Short day induced bud break, due to an activated rib meristem capable of inducing callose hydrolysis at plasmodesmata, was later linked to increased GA level, synthesised from an abundant pool of

precursors within the dwarf shoot (Rinne et al. 2016). Thus, the physical properties of the rib meristem may interact with its responsiveness and capacity to elongate which may be relevant to intercalary meristem activation timing and pattern along the stem. Intriguingly, the *Carbohydrate partitioning defective1* dominant mutant in maize is developmentally delayed, dwarf and shows excessive phloem callose plugs and lignification, associated with inability to mobilise starch out of the leaves (Julius et al. 2018). It will be interesting to learn how callose deposition and sugar movement (see below) may relate to intercalary meristem dormancy/ activation.

Transport in and out of the intercalary meristem also influences its activity. The maize *brachytic2* (*br2*) and sorghum *dwarf3* mutants have short, thick internodes caused by defects in a gene encoding an ATP-binding cassette type B1 (ABCB1) auxin efflux transporter (Multani et al. 2003). Knöller et al (2010) showed that *BR2* was expressed in the intercalary meristem and that *br2* mutants showed abnormal nodal vasculature, defective auxin transport out of the intercalary meristem and reduced intercalary proliferation. Thus, proliferative activity in intercalary meristems is likely dependent on proper nodal vasculature supporting auxin export, an intriguing parallel to the role for polar auxin transport out of the bud in promoting axillary bud outgrowth versus dormancy (Domagalska and Leyser 2011). In the case of the axillary bud, this is due to PIN localisation dynamics in the stem and bud vasculature (Balla et al. 2011; Domagalska and Leyser 2011; Shinohara et al. 2013).

Determining vascular patterning, auxin fluxes and PIN dynamics in intercalary meristems both before and during activation may be important to understand the relevance of auxin to internode elongation, and how these mechanisms interact with the aforementioned KNOX and BLH functions.

Internodes are adjacent to and develop from same the phytomer as axillary buds. An inverse relationship between internode elongation and axillary bud outgrowth is often observed in cereals. Bud outgrowth in cereals, called tillering, occurs from nodes between basal unexpanded internodes and stops when internode elongation starts (Kirby and Appleyard 1984). Thus, expanding internodes generally do not have tillers, although this relationship is broken in GA constitutive mutants in barley where basal internode elongation occurs concurrently with tillering, and plants adopt an aerial branching habit (Foster 1977), as is also observed in the many-noded dwarf mutants (Harlan and Pope 1922). Another consideration is why tiller buds adjacent to elongating internodes do not grow out. In his recent, intriguing review, Kebrom (2017) postulates that the increased sugar demand from expanding internodes may divert sugar away from their axillary buds, thus promoting dominance of the elongating shoot. Each subsequent internode tends to be longer than its underlying internode (Briggs 1978): i.e. Internodes increase in length with further distance from the outgrowing tillers, potentially

reflecting this interaction. Furthermore, given the role for auxin export from the axillary bud, and a potential role for this movement in intercalary meristem activity (Knöller et al. 2011), the intercalary meristem and the axillary bud rib meristem could also compete for auxin export. Interestingly, strigalactone (SL), which inhibits axillary bud growth by influencing PIN-mediated auxin efflux out of the bud, also promotes internode elongation (de Saint Germain et al. 2013).

CONCLUSION AND OUTLOOK

The Green Revolution heralded the most spectacular increase in food supply in human history (<https://ourworldindata.org/>). Selection for variation in growth hormone pathways controlling stem elongation was critical to increase yield, yet also brought some undesirable pleiotropic effects throughout the plant lifecycle, such as reduced seed germination and vigour in GA signalling mutants (Wu et al. 2011; Nagel et al. 2013). Although we know an impressive amount about the downstream molecular mechanisms of these hormones, we are less sure about other components, potentially upstream, which may specifically regulate both primary stem morphogenesis and the secondary stem elongation so critical to yield. Understanding these regulators may help not only selectively control internode growth compared to other features but gain insight about the pathways controlling upward plant growth, a fundamental activity with profound implications on plant development, diversity and evolution.

Several questions to pursue include:

- 1) Are there CLV-WUS/KNOX meristem maintenance pathways specific for the rib meristem and/ or intercalary meristem maintenance?
- 2) Much of our understanding about stem growth is from work in Arabidopsis, pea and the warm weather rice and maize cereals. How conserved are these mechanisms in temperate cereals?
- 3) How conserved are regulators of intercalary or rib meristem activity compared to other meristems in non-vascular plants which give rise to transverse cell files during sporophyte elevation? How does this reflect the present/absence of lignin between these two groups?
- 4) Is there an apically-derived signal activating intercalary activity and how is it perceived?
- 5) How is vascular differentiation in the node, internode and axillary bud regulated and is this relevant to their differential pattern of internode growth within an individual? In particular, how is this timed and relevant to rib meristem function and intercalary dormancy?
- 6) How are environmental signals perceived at either the rib or intercalary meristem to alter their activity in response to changed conditions during

stem elongation and are these signals more important for proliferation or expansion in the internode?

ACKNOWLEDGEMENTS

I acknowledge support from a Royal Society of Edinburgh Personal Research Fellowship and from a BBSRC New Investigator Grant (No. BB/L001934/1).

REFERENCES

- Adibi M, Yoshida S, Weijers D, Fleck C (2016) Centering the organizing center in the *Arabidopsis thaliana* shoot apical meristem by a combination of cytokinin signaling and self-organization. **PLoS ONE** 11: e0154638
- Aguilar-Martinez JA, Poza-Carrion C, Cubas P (2007) *Arabidopsis* BRANCHED1 acts as an integrator of branching signals within axillary buds **Plant Cell** 19: 458–472
- Angeles-Shim RB, Asano K, Takashi T, Shim J, Kuroha T, Ayano M, Ashikari M (2012) A WUSCHEL-related homeobox 3B gene, *depilous (dep)*, confers glabrousness of rice leaves and glumes. **Rice** 5: 1–12
- Aoki T, Kitano H, Kameya N, Nakamura I (2002) Accelerated shoot overgrowth of rice mutant *ao-1* is epistatic to gibberellin-sensitive and -insensitive dwarf mutants. **J Plant Res** 115: 195–202
- Ashikari M, Wu J, Yano M, Sasaki T, Yoshimura A (1999) Rice gibberellin-insensitive dwarf mutant gene Dwarf 1 encodes the α -subunit of GTP-binding protein. **Proc Natl Acad Sci USA** 96: 10284–10289
- Bai MY, Shang JX, Oh E, Fan M, Bai Y, Zentella R, Sun TP, Wang ZY (2012) Brassinosteroid, gibberellin and phytochrome impinge on a common transcription module in *Arabidopsis*. **Nat Cell Biol** 14: 810–817
- Baima S, Forte V, Possenti M, Peñalosa A, Leoni G, Salvi S, Felici B, Ruberti I, Morelli G (2014) Negative feedback regulation of auxin signaling by ATHB8/ACL5–BUD2 transcription module. **Mol Plant** 7: 1006–1025
- Baima S, Nobili F, Sessa G, Lucchetti S, Ruberti I, Morelli G (1995) The expression of the *Athb-8* homeobox gene is restricted to provascular cells in *Arabidopsis thaliana*. **Development** 121: 4171 LP-4182
- Baima S, Possenti M, Matteucci A, Wisman E, Altamura MM, Ruberti I, Morelli G (2001) The *Arabidopsis* ATHB-8 HD-Zip protein acts as a differentiation-promoting transcription factor of the vascular meristems. **Plant Physiol** 126: 643–655
- Balla J, Kalousek P, Reinöhl V, Friml J, Procházka S (2011) Competitive canalization of PIN-dependent auxin flow from axillary buds controls pea bud

outgrowth. **Plant J** 65: 571–577

Becraft PW, Freeling M (1994) Genetic analysis of Rough sheath1 developmental mutants of maize. **Genetics** 136: 295 LP-311

Bencivenga S, Serrano-Mislata A, Bush M, Fox S, Sablowski R (2016) Control of oriented tissue growth through repression of organ boundary genes promotes stem morphogenesis. **Dev Cell** 39: 198–208

Binenbaum J, Weinstein R, Shani E (2018) Gibberellin localization and transport in plants. **Trends Plant Sci** 23:410–421

Bleecker AB, Schuette JL, Kende H (1986) Anatomical analysis of growth and developmental patterns in the internode of deepwater rice. **Planta** 169: 490–497

Bolduc N, Hake S (2009) The maize transcription factor KNOTTED1 directly regulates the gibberellin catabolism Gene *ga2ox1*. **Plant Cell** 21: 1647–1658

Bommert P, Je B II, Goldshmidt A, Jackson D (2013) The maize Gα gene *COMPACT PLANT2* functions in CLAVATA signalling to control shoot meristem size. **Nature** 502: 555–558

Bommert P, Lunde C, Nardmann J, Vollbrecht E, Running MP, Jackson D, Hake S, Werr W (2005) thick tassel dwarf1 encodes a putative maize ortholog of the *Arabidopsis* CLAVATA1 leucine-rich repeat receptor-like kinase. **Development** 132: 1235–1245

Bommert P, Whipple C (2018) Grass inflorescence architecture and meristem determinacy. *Semin. Cell Dev Biol* 79: 37–47

Bossinger G (2009) Segments (Phytomers). eLS

Brand U, Grünwald M, Hobe M, Simon R (2002) Regulation of *CLV3* expression by two homeobox genes in *Arabidopsis*. **Plant Physiol** 129: 565–575

Braumann I, Dockter C, Beier S, Himmelbach A, Lok F, Lundqvist U, Skadhauge B, Stein N, Zakhrebekova S, Zhou R, Hansson M (2018) Mutations in the gene of the Gα subunit of the heterotrimeric G protein are the cause for the *brachytic1* semi-dwarf phenotype in barley and applicable for practical breeding. **Hereditas** 155: 10

Briggs DE (1978) Barley. Chapman & Hall, London

Byrne ME, Simorowski J, Martienssen RA (2002) *ASYMMETRIC LEAVES1* reveals *knox* gene redundancy in *Arabidopsis*. **Development** 129: 1957–1965

Chickarmane VS, Gordon SP, Tarr PT, Heisler MG, Meyerowitz EM (2012) Cytokinin signaling as a positional cue for patterning the apical-basal axis of the growing *Arabidopsis* shoot meristem. **Proc Natl Acad Sci USA** 109: 4002–4007

Cho H, Kende H (1997) Expansins in deepwater rice internodes. **Plant Physiol** 113: 1137–1143

Cho S-H, Kang K, Lee S-H, Lee I-J, Paek N-C (2016) OsWOX3A is involved in negative feedback regulation of the gibberellic acid biosynthetic pathway in rice (*Oryza sativa*). **J Exp Bot** 67: 1677–87

Cho SH, Yoo SC, Zhang H, Pandeya D, Koh HJ, Hwang JY, Kim GT, Paek NC (2013) The rice *narrow leaf2* and *narrow leaf3* loci encode WUSCHEL-related homeobox 3A (OsWOX3A) and function in leaf, spikelet, tiller and lateral root development. **New Phytol** 198: 1071–1084

Chono M, Honda I, Zeniya H, Yoneyama K, Saisho D, Takeda K, Takatsuto S, Hoshino T, Watanabe Y (2003) A semidwarf phenotype of barley uzu results from a nucleotide substitution in the gene encoding a putative brassinosteroid receptor. **Plant Physiol** 133: 1209–1219

Clark S, Running M, Meyerowitz E (1995) *CLAVATA3* is a specific regulator of shoot and floral meristem development affecting the same processes as *CLAVATA1*. **Development** 121: 2057–2067

Clark SE, Jacobsen SE, Levin JZ, Meyerowitz EM (1996) The *CLAVATA* and *SHOOT MERISTEMLESS* loci competitively regulate meristem activity in *Arabidopsis*. **Development** 122: 1567–75

Clark SE, Running MP, Meyerowitz EM (1993) *CLAVATA1*, a regulator of meristem and organ development in *Arabidopsis*. **Development** 125: 3843–3851

Clay NK, Nelson T (2005) *Arabidopsis* thickvein mutation affects vein thickness and organ vascularization, and resides in a provascular cell-specific spermine synthase involved in vein definition and in polar auxin transport. **Plant Physiol** 138: 767–777

Cole M, Nolte C, Werr W (2006) Nuclear import of the transcription factor SHOOT MERISTEMLESS depends on heterodimerization with BLH proteins expressed in discrete sub-domains of the shoot apical meristem of *Arabidopsis thaliana*. **Nucleic Acids Res** 34: 1281–1292

Cronk Q (2009) The Molecular Organography of Plants. Oxford University Press, oxford

Cui K, He CY, Zhang JG, Duan AG, Zeng YF (2012) Temporal and spatial profiling of internode elongation-associated protein expression in rapidly growing culms of bamboo. **J Proteome Res** 11: 2492–2507

Dai M, Hu Y, Ma Q, Zhao Y, Zhou D-X (2008) Functional analysis of rice *HOMEBOX4* (*Oshox4*) gene reveals a negative function in gibberellin responses. **Plant Mol Biol** 66: 289–301

Davière J-M, Achard P (2013) Gibberellin signaling in plants. **Development** 140: 1147–1151

Davière J-M, Wild M, Regnault T, Baumberger N, Eisler H, Genschik P, Achard P (2014) Class I TCP-DELLA interactions in inflorescence shoot apex determine plant height. **Curr Biol** 24: 1923–1928

Dayan J, Voronin N, Gong F, Sun T, Hedden P, Fromm H, Aloni R (2012) Leaf-induced gibberellin signaling is essential for internode elongation, cambial activity, and fiber differentiation in tobacco stems. **Plant Cell** 24: 66–79

de Saint Germain A, Ligerot Y, Dun EA, Pillot J-P, Ross JJ, Beveridge CA, Rameau C (2013) Strigolactones stimulate internode elongation independently of gibberellins. **Plant Physiol** 163: 1012–1025

DeYoung BJ, Clark S (2008) BAM receptors regulate stem cell specification and organ development through complex interactions with CLAVATA signaling. **Genetics** 180: 895–904

Dockter C, Gruszka D, Braumann I, Druka A, Druka I, Franckowiak J, Gough SP, Janeczko A, Kurowska M, Lundqvist J, Lundqvist U, Marzec M, Matyszcak I, Müller AH, Oklestkova J, Schulz B, Zakhrebekova S, Hansson M (2014) Induced variations in brassinosteroid genes define barley height and sturdiness, and expand the green revolution genetic toolkit. **Plant Physiol** 166: 1912–1927

Dockter C, Hansson M (2015) Improving barley culm robustness for secured crop yield in a changing climate. **J Exp Bot** 66: 3499–3509

Domagalska MA, Leyser O (2011) Signal integration in the control of shoot branching. **Nat Rev Mol Cell Biol** 12: 211–221

Ehlting J1, Mattheus N, Aeschliman DS, Li E, Hamberger B, Cullis IF, Zhuang J, Kaneda M, Mansfield SD, Samuels L, Ritland K, Ellis BE, Bohlmann J, Douglas CJ (2005) Global transcript profiling of primary stems from *Arabidopsis thaliana* identifies candidate genes for missing links in lignin biosynthesis and transcriptional regulators of fiber differentiation. **Plant J** 42: 618–640

Emery JF, Floyd SK, Alvarez J, Eshed Y, Hawker NP, Izhaki A, Baum SF, Bowman JL (2003) Radial patterning of *Arabidopsis* shoots by class III HD-ZIP and KANADI genes. **Curr Biol** 13: 1768–1774

Eriksson S, Böhlenius H, Moritz T, Nilsson O (2006) GA4 is the active gibberellin in the regulation of *LEAFY* transcription and *Arabidopsis* floral initiation. **Plant Cell** 18: 2172–2181

Etchells JP, Provost CM, Mishra L, Turner SR (2013) *WOX4* and *WOX14* act downstream of the PXY receptor kinase to regulate plant vascular proliferation independently of any role in vascular organisation. **Development** 140: 2224–2234

Etchells JP, Turner SR (2010) The PXY-CLE41 receptor ligand pair defines a multifunctional pathway that controls the rate and orientation of vascular cell

division. **Development** 137: 767–774

Evans PS (1965) Intercalary growth in the aerial shoot of *eleocharis acuta* R. Br. Prodr.: I. structure of the growing zone 1. **Ann Bot** 29: 205–217

Fahn A, Stoler S, First T (1963) Vegetative shoot apex in banana and zona I changes as it becomes reproductive. **Bot Gaz** 124: 246–250

Fambrini M, Bonsignori E, Rapparini F, Cionini G, Michelotti V, Bertini D, Baraldi R, Pugliesi C (2006) Stem fasciated, a recessive mutation in sunflower (*Helianthus annuus*), alters plant morphology and auxin level. **Ann Bot** 98: 715–730

Fisher JB (1970) Development of the intercalary meristem of *cyperus alternifolius*. **Am J Bot** 57: 691–703

Fisher JB, French JC (1976) The occurrence of intercalary and uninterrupted meristems in the internodes of tropical monocotyledons. **Am J Bot** 63: 510

Fisher JB, French JC (1978) Internodal meristems of monocotyledons: Further studies and a general taxonomic survey. **Ann Bot** 42: 41–50

Fletcher JC (2002) Coordination of cell proliferation and cell fate decisions in the angiosperm shoot apical meristem. **Bioessays** 24: 27–37

Foster CA (1977) Slender: An accelerated extension growth mutant of barley. **Barley Genet Newsl** 7: 24–27

Foster T, Yamaguchi J, Wong BC, Veit B, Hake S (1999) *Gnarley1* is a dominant mutation in the *knox4* Homeobox gene affecting cell shape and identity. **Plant Cell** 11: 1239–1252

Freeling M, Hake S (1985) Developmental genetics of mutants that specify knotted leaves in maize. **Genetics** 111: 617–634

Fujisawa Y, Kato T, Ohki S, Ishikawa A, Kitano H, Sasaki T, Asahi T, Iwasaki Y (1999) Suppression of the heterotrimeric G protein causes abnormal morphology, including dwarfism, in rice. **Proc Natl Acad Sci USA** 96: 7575–7580

Gaillochet C, Daum G, Lohmann JU (2015) O cell, where art thou? The mechanisms of shoot meristem patterning. **Curr Opin Plant Biol** 23: 91–97

Gaillochet C, Jamge S, van der Wal F, Angenent G, Immink R, Lohmann JU (2018) A molecular network for functional versatility of HECATE transcription factors. **Plant J** 95: 57–70

Gale M, Youssefian S (1985) Dwarfing genes in wheat. In: Russell G, ed. *Progress in Plant Breeding*. Butterworth, London. pp. 1–35

Galinat WC (1959) The phytomer in relation to floral homologies in the Amercian Maydeae. **Bot Mus Lealf Harv Univ** 19: 1–xvi

- Gallego-Bartolomé J, Minguet EG, Grau-Enguix F, Abbas M, Locascio A, Thomas SG, Alabadí D, Blázquez MA (2012) Molecular mechanism for the interaction between gibberellin and brassinosteroid signaling pathways in *Arabidopsis*. **Proc Natl Acad Sci USA** 109: 13446–51
- Gao S, Fang J, Xu F, Wang W, Chu C (2016) Rice HOX12 regulates panicle exertion by directly modulating the expression of *ELONGATED UPPERMOST INTERNODE1*. **Plant Cell** 28: 680–695
- Gómez-Mena C, Sablowski R (2008) *ARABIDOPSIS THALIANA* *ARABIDOPSIS THALIANA* HOMEBOX GENE1 establishes the basal boundaries of shoot organs and controls stem growth. **Plant Cell** 20: 2059–2072
- Gordon S, Chickarmane V (2009) Multiple feedback loops through cytokinin signaling control stem cell number within the *Arabidopsis* shoot meristem. **Proc Natl Acad Sci USA** 106: 16529–16534
- Gorter CJ (1965) Origin of fasciation BT - Differenzierung und Entwicklung In: Lang A, ed. *Differentiation and Development*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 1977–1998
- Greb T, Lohmann JU (2016) Plant stem cells. **Curr Biol** 26: R816–R821
- Gregory EF, Dao TQ, Alexander MA, Miller MJ, Fletcher JC (2018) The signaling peptide-encoding genes *CLE16*, *CLE17* and *CLE27* are dispensable for *Arabidopsis* shoot apical meristem activity. **PLoS ONE** 13: e0202595
- Grisebach A (1943) Beobachtungen fiber das Wachstum der Vegetationsorgane in Bezug auf Systemati. **Arch Naturgesch** 9: 267–292
- Gruel J, Landrein B, Tarr P, Schuster C, Refahi Y, Sampathkumar A, Hamant O, Meyerowitz EM, Jönsson H (2016) An epidermis-driven mechanism positions and scales stem cell niches in plants. **Sci Adv** 2: e1500989
- Harlan H V, Pope MN (1922) Many-noded dwarf barley. **J Hered** 13: 269–273
- Harrison CJ, Corley SB, Moylan EC, Alexander DL, Scotland RW, Langdale JA (2005) Independent recruitment of a conserved developmental mechanism during leaf evolution. **Nature** 434: 509
- Harrison CJ, Morris JL (2018) The origin and early evolution of vascular plant shoots and leaves. **Philos Trans R Soc Lond B Biol Sci** 373: 20160496
- Harting M (1845) Recherches micrométriques sur le développement des parties élémentaires de la tige annuelle des plantes dicotylédonées. **Ann Sci NatBot Ser III** 4: 210–279
- Hay A, Kaur H, Phillips A, Hedden P, Hake S, Tsiantis M (2002) The gibberellin pathway mediates KNOTTED1-Type homeobox function in plants with different body plans. **Curr Biol** 12: 1557–1565

Hedden P (2003) The genes of the Green Revolution. **Trends Genet** 19: 5–9

Hempel FD, Feldman LJ (1994) Bi-directional inflorescence development in *Arabidopsis thaliana*: Acropetal initiation of flowers and basipetal initiation of bracteoles. **Planta** 192: 276–286

Huang X, Qian Q, Liu Z, Sun H, He S, Luo D, Xia G, Chu C, Li J, Fu X (2009) Natural variation at the *DEP1* locus enhances grain yield in rice. **Nat Genet** 41: 494

Hubbard L, McSteen P, Doebley J, Hake S (2002) Expression patterns and mutant phenotype of teosinte branched1 correlate with growth suppression in maize and teosinte. **Genetics** 162: 1927–1935

Ikeda A, Ueguchi-Tanaka M, Sonoda Y, Kitano H, Koshioka M, Futsuhara Y, Matsuoka M, Yamaguchi J (2001) *slender rice*, a constitutive gibberellin response mutant, is caused by a null mutation of the *SLR1* gene, an ortholog of the height-regulating gene *GAI/RGA/RHT/D8*. **Plant Cell** 13: 999–1010

Ishida T, Tabata R, Yamada M, Aida M, Mitsumasu K, Fujiwara M, Yamaguchi K, Shigenobu S, Higuchi M, Tsuji H, et al (2016) Heterotrimeric G proteins control stem cell proliferation through CLAVATA signaling in *Arabidopsis*. **EMBO Rep** 17: 1236–1236

Ishiwata A, Ozawa M, Nagasaki H, Kato M, Noda Y, Yamaguchi T, Nosaka M, Shimizu-Sato S, Nagasaki A, Maekawa M, et al (2013) Two WUSCHEL-related homeobox genes, *narrow leaf2* and *narrow leaf3*, control leaf width in rice. **Plant Cell Physiol** 54: 779–792

Ito A, Yasuda A, Yamaoka K, Ueda M, Nakayama A, Takatsuto S, Honda I (2017) Brachytic 1 of barley (*Hordeum vulgare* L.) encodes the alpha subunit of heterotrimeric G protein. **J Plant Physiol** 213: 209–215

Jackson D, Veit B, Hake S (1994) Expression of maize *KNOTTED1* related homeobox genes in the shoot apical meristem predicts patterns of morphogenesis in the vegetative shoot. **Development** 121: 405–413

Jacqumard A, Gadisseur I, Bernier G (2003) Cell division and morphological changes in the shoot apex of *Arabidopsis thaliana* during floral transition. **Ann Bot** 91: 571–576

Janocha D, Lohmann JU (2018) From signals to stem cells and back again. **Curr Opin Plant Biol** 45: 136–142

Jasinski S, Piazza P, Craft J, Hay A, Woolley L, Rieu I, Phillips A, Hedden P, Tsiantis M (2005) KNOX action in *Arabidopsis* is mediated by coordinate regulation of cytokinin and gibberellin activities. **Curr Biol** 15: 1560–1565

Je B II, Gruel J, Lee YK, Bommert P, Arevalo ED, Eveland AL, Wu Q, Goldshmidt A, Meeley R, Bartlett M, et al (2016) Signaling from maize organ primordia via FASCIATED EAR3 regulates stem cell proliferation and yield traits. **Nat Genet** 48: 785–791

Jegla DE, Sussex IM (1989) Cell lineage patterns in the shoot meristem of the sunflower embryo in the dry seed. **Dev Biol** 131: 215–225

Johri MM, Coe EH (1983) Clonal analysis of corn plant development. I. The development of the tassel and the ear shoot. **Dev Biol** 97: 154–172

Julius BT, Slewinski TL, Baker RF, Tzin V, Zhou S, Bihmidine S, Jander G, Braun DM (2018) Maize Carbohydrate partitioning defective1 impacts carbohydrate distribution, callose accumulation, and phloem function. **J Exp Bot** 69: 3917–3931

Jun J, Fiume E, Roeder AHK, Meng L, Sharma VK, Osmont KS, Baker C, Ha CM, Meyerowitz EM, Feldman LJ, et al (2010) Comprehensive analysis of *CLE* polypeptide signaling gene Expression and overexpression activity in *Arabidopsis*. **Plant Physiol** 154: 1721–1736

Takehi J, Kuwashiro Y, Niitsu M, Takahashi T (2008) Thermospermine is required for stem elongation in *Arabidopsis thaliana*. **Plant Cell Physiol** 49: 1342–1349

Kaufman PB (1959) Development of the shoot of *Oryza sativa* L. III. Early stages in histogenesis of the stem and ontogeny of the adventitious root. **Phytomorph** 9: 382–404

Kaufman PB, Cassell SJ, Adams PA (1965) On nature of intercalary growth and cellular differentiation in internodes of *Avena sativa*. **Bot Gaz** 126: 1–13

Kaya H, Shibahara K ichi, Taoka K ichiro, Iwabuchi M, Stillman B, Araki T (2001) *FASCIATA* genes for chromatin assembly factor-1 in *Arabidopsis* maintain the cellular organization of apical meristems. **Cell** 104: 131–142

Kayes JM, Clark SE (1998) *CLAVATA1*, a regulator of meristem and organ development in *Arabidopsis*. **Development** 125: 3843–3851

Kebrom TH (2017) A growing stem inhibits bud outgrowth – The overlooked theory of apical dominance. **Front Plant Sci** 8: 1874

Kebrom TH, McKinley B, Mullet JE (2017) Dynamics of gene expression during development and expansion of vegetative stem internodes of bioenergy sorghum. **Biotechnol Biofuels** 10: 159

Kellogg EA (2015) Flowering Plants. Monocots. Springer International Publishing AG, Cham

Kende H, van der Knaap E E van der, Cho HT (1998) Deepwater rice: A model plant to study stem elongation. **Plant Physiol** 118: 1105–10

Khan M, Xu M, Murmu J, Tabb P, Liu Y, Storey K, McKim SM, Douglas CJ, Hepworth SR (2012) Antagonistic interaction of *BLADE-ON-PETIOLE1* and 2 with *BREVIPEDICELLUS* and *PENNYWISE* regulates *Arabidopsis* inflorescence architecture. **Plant Physiol** 158: 946–960

Kieffer M, Master V, Waites R, Davies B (2011) TCP14 and TCP15 affect internode length and leaf shape in *Arabidopsis*. **Plant J** 68: 147–158

Kirby EJM, Appleyard M (1984) Cereal development guide. 2nd Edition. Arable Unit, National Agricultural Centre, Stoneleigh, Warwickshire

Kirby EJM, Appleyard M, Simpson NA (1994) Co-ordination of stem elongation and Zadoks growth stages with leaf emergence in wheat and barley. **J Agric Sci** 122: 21-29

Knöller AS, Blakeslee JJ, Richards EL, Peer WA, Murphy AS (2010) Brachytic2/ZmABCB1 functions in IAA export from intercalary meristems. **J Exp Bot** 61: 3689–3696

Koning R, Tkaczyk A, Kaufman PB, Pharis RP, Morf W (1977) Regulation of internodal extension in *Avena* shoots by the inflorescence, nodes, leaves, and intercalary meristem. **Physiol Plant** 40: 119–124

Kwiatkowska D (2008) Flowering and apical meristem growth dynamics. **J Exp Bot** 59: 187–201

Laux T, Mayer KF, Berger J, Jürgens G (1996) The *WUSCHEL* gene is required for shoot and floral meristem integrity in *Arabidopsis*. **Development** 122: 87–96

Leibfried A, To JPC, Busch W, Stehling S, Kehle A, Demar M, Kieber JJ, Lohmann JU (2005) WUSCHEL controls meristem function by direct regulation of cytokinin-inducible response regulators. **Nature** 438: 1172–1175

Leyser OH, Furner I (1992) Characterisation of three shoot apical meristem mutants of *Arabidopsis thaliana*. **Development** 116: 397–403

Li Q-F, Wang C, Jiang L, Li S, Sun SSM, He J-X (2012) An interaction between BZR1 and DELLAs mediates direct signaling crosstalk between brassinosteroids and gibberellins in *Arabidopsis*. **Sci Signal** 5: ra72

Ligrone R, Duckett JG, Renzaglia KS (2012) Major transitions in the evolution of early land plants: A bryological perspective. **Ann Bot** 109: 851–871

Lincoln C, Long J, Yamaguchi J, Serikawa K, Hake S (1994) A *knotted1*-like homeobox gene in *Arabidopsis* is expressed in the vegetative meristem and dramatically alters leaf morphology when overexpressed in transgenic plants. **Plant Cell** 6: 1859-1876

Liu C, Zheng S, Gui J, Fu C, Yu H, Song D, Shen J, Qin P, Liu X, Han B, et al (2018) *Shortened Basal Internodes* encodes a gibberellin 2-oxidase and contributes to lodging resistance in rice. **Mol Plant** 11: 288–299

Mandel T, Candela H, Landau U, Asis L, Zelinger E, Carles CC, Williams LE (2016) Differential regulation of meristem size, morphology and organization by the ERECTA, CLAVATA and class III HD-ZIP pathways. **Development** 143: 1612 LP-1622

Martín-Trillo M, Cubas P (2010) *TCP* genes: A family snapshot ten years later. **Trends Plant Sci** 15: 31–9

Martin AP, Palmer WM, Brown C, Abel C, Lunn JE, Furbank RT, Grof CPL (2016) A developing *Setaria viridis* internode: An experimental system for the study of biomass generation in a C₄ model species. **Biotechnol Biofuels** 9: 45

Matsumoto N, Okada K (2001) A homeobox gene, *PRESSED FLOWER*, regulates lateral axis-dependent development of *Arabidopsis* flowers. **Genes Dev** 15: 3355–3364

McDaniel CN, Poethig RS (1988) Cell-lineage patterns in the shoot apical meristem of the germinating maize embryo. **Planta** 175: 13–22

Medford J, Behringer F, Callos J, Feldmann K (1992) Normal and abnormal development in the *Arabidopsis* vegetative shoot apex. **Plant Cell** 4: 631–643

Mele G, Ori N, Sato Y, Hake S (2003) The knotted1-like homeobox gene *BREVIPEDICELLUS* regulates cell differentiation by modulating metabolic pathways. **Genes Dev** 17: 2088–2093

Meng W, Cheng ZJ, Sang YL, Zhang MM, Rong XF, Wang ZW, Tang YY, Zhang XS (2017) Type-B ARABIDOPSIS RESPONSE REGULATORS specify the shoot stem cell Niche by dual regulation of *WUSCHEL*. **Plant Cell** 29: 1357–1372

Moon S, Jung KH, Lee DE, Lee DY, Lee J, An K, Kang HG, An G (2006) The rice *FON1* gene controls vegetative and reproductive development by regulating shoot apical meristem size. **Mol Cells** 21: 147–152

Muller R, Borghi L, Kwiatkowska D, Laufs P, Simon R (2006) Dynamic and compensatory responses of *Arabidopsis* shoot and floral meristems to CLV3 signaling. **Plant Cell** 18: 1188–1198

Multani DS, Briggs SP, Chamberlin MA, Blakeslee JJ, Murphy AS, Johal GS (2003) Loss of an MDR transporter in compact stalks of maize *br2* and sorghum *dw3* mutants. **Science** 302: 81–84

Muñoz-Nortes T, Wilson-Sánchez D, Candela H, Micol JL (2014) Symmetry, asymmetry, and the cell cycle in plants: Known knowns and some known unknowns. **J Exp Bot** 65: 2645–2655

Nagel M, Behrens A-K, Börner A (2013) Effects of *Rht* dwarfing alleles on wheat seed vigour after controlled deterioration. **Crop Pasture Sci** 64: 857–864

Nardmann J, Ji J, Werr W, Scanlon MJ (2004) The maize duplicate genes *narrow sheath1* and *narrow sheath2* encode a conserved homeobox gene function in a lateral domain of shoot apical meristems. **Development** 131: 2827–2839

Nardmann J, Werr W (2006) The shoot stem cell niche in angiosperms: Expression patterns of WUS orthologues in rice and maize imply major modifications in the course of mono- and dicot evolution. **Mol Biol Evol** 23: 2492–2504

Nardmann J, Zimmermann R, Durantini D, Kranz E, Werr W (2007) *WOX* gene phylogeny in poaceae: A comparative approach addressing leaf and embryo development. **Mol Biol Evol** 24: 2474–2484

Nicholls PB, May LH (1964) Studies on the growth of the barley apex. II. on the initiation of internode elongation in the inflorescence. **Aust J Bio Sci** 17: 619–630

Niklas KJ (1990) The mechanical significance of clasping leaf sheaths in grasses: Evidence from two cultivars of *Avena sativa*. **Ann Bot** 65: 505–512

Nishikawa N, Toyama S, Shida A, Futatsuya F (1994) The uptake and the transport of ¹⁴C-labeled epibrassinolide in intact seedlings of cucumber and wheat. **J Plant Res** 107: 125–130

Norberg M, Holmlund M, Nilsson O (2005) The *BLADE ON PETIOLE* genes act redundantly to control the growth and development of lateral organs. **Development** 132: 2203–2213

Ohmori Y, Tanaka W, Kojima M, Sakakibara H, Hirano H-Y (2013) *WUSCHEL-RELATED HOMEODOMAIN 4* Is Involved in Meristem Maintenance and Is Negatively Regulated by the CLE Gene *FCP1* in Rice. **Plant Cell** 25: 229–241

Oki K, Inaba N, Kitano H, Takahashi S, Fujisawa Y, Kato H, Iwasaki Y (2009) Study of novel *d1* alleles, defective mutants of the alpha subunit of heterotrimeric G-protein in rice. **Genes Genet Syst** 84: 35–42

Pautler M, Eveland AL, LaRue T, Yang F, Weeks R, Lunde C, Je B II, Meeley R, Komatsu M, Vollbrecht E, Sakai H, Jackson D (2015) *FASCIATED EAR4* encodes a bZIP transcription factor that regulates shoot meristem size in maize. **Plant Cell** 27: 104–120

Pautler M, Tanaka W, Hirano HY, Jackson D (2013) Grass meristems I: shoot apical meristem maintenance, axillary meristem determinacy and the floral transition. **Plant Cell Physiol** 54: 302–12

Peng J, Carol P, Richards DE, King KE, Cowling RJ, Murphy GP, Harberd NP (1997) The *Arabidopsis* *GAI* gene defines a signaling pathway that negatively regulates gibberellin responses. **Genes Dev** 11: 3194–3205

Peng J, Richards DE, Hartley NM, Murphy GP, Devos KM, Flintham JE, Beales J, Fish LJ, Worland AJ, Pelica F, Sudhakar D, Christou P, Snape JW, Gale MD, Harberd NP (1999) “Green revolution” genes encode mutant gibberellin response modulators. **Nature** 400: 256–261

Perales M, Reddy GV (2012) Stem cell maintenance in shoot apical

meristems. **Curr Opin Plant Biol** 15: 10–6

Poethig RS, Szymkowiak EJ (1995) Clonal analysis of leaf development in maize. **Maydica** 40: 67–76

Pouteau S, Albertini C (2011) An assessment of morphogenetic fluctuation during reproductive phase change in *Arabidopsis*. **Ann Bot** 107: 1017–1027

Prigge MJ, Otsuga D, Alonso JM, Ecker JR, Drews GN, Clark SE (2005) Class III homeodomain-leucine zipper gene family members have overlapping, antagonistic, and distinct roles in *Arabidopsis* development. **Plant Cell** 17: 61–76

Ragni L, Belles-Boix E, Günl M, Pautot V (2008) Interaction of *KNAT6* and *KNAT2* with *BREVIPEDICELLUS* and *PENNYWISE* in *Arabidopsis* inflorescences. **Plant Cell** 20: 888–900

Reiser L, Sanchez-Baracaldo P, Hake S (2000) Knots in the family tree: Evolutionary relationships and functions of *knox* homeobox genes. **Plant Mol Biol** 42: 151–166

Rinne PLH, Paul LK, Vahala J, Kangasjärvi J, Van Der Schoot C (2016) Axillary buds are dwarfed shoots that tightly regulate GA pathway and GA-inducible 1,3- β -glucanase genes during branching in hybrid aspen. **J Exp Bot** 67: 5975–5991

Ross JJ, O'Neill DP, Smith JJ, Kerckhoffs LHJ, Elliott RC (2000) Evidence that auxin promotes gibberellin A1 biosynthesis in pea. **Plant J** 21: 547–552

Running MP, Lavy M, Sternberg H, Galichet A, Gruissem W, Hake S, Ori N, Yalovsky S (2004) Enlarged meristems and delayed growth in *plp* mutants result from lack of CaaX prenyltransferases. **Proc Natl Acad Sci USA** 101: 7815–7820

Ruonala R, Rinne PLH, Kangasjarvi J, van der Schoot C (2008) *CENL1* expression in the rib meristem affects stem elongation and the transition to dormancy in *Populus*. **Plant Cell** 20: 59–74

Sachs RM (1965) Stem elongation. **Annu Rev Plant Physiol** 16: 73–96

Sachs RM, Bretz C, Lang A (1959a) Cell division and gibberellic acid. **Exp Cell Res** 18: 230–244

Sachs RM, Bretz CF, Lang A (1959b) Shoot histogenesis: The early effects of gibberellin upon stem elongation in two rosette plants. **Am J Bot** 46: 376

Sachs RM, Lang A, Bretz CF, Roach J (1960) Shoot histogenesis: Subapical meristematic activity in a caulescent plant and the action of gibberellic acid and Amo-1618. **Am J Bot** 47: 260–266

Sanchez P, Nehlin L, Greb T (2012) From thin to thick: Major transitions during stem development. **Trends Plant Sci** 17: 113–21

Sato Y, Sentoku N, Miura Y, Hirochika H, Kitano H, Matsuoka M (1999) Loss-of-function mutations in the rice homeobox gene OSH15 affect the architecture of internodes resulting in dwarf plants. **EMBO J** 18: 992–1002

Scanlon MJ (2003) The polar auxin transport inhibitor *N*-1-naphthylphthalamic acid disrupts leaf initiation, KNOX protein regulation, and formation of leaf margins in maize. **Plant Physiol** 133: 597–605

Scanlon MJ, Freeling M (1997) Clonal sectors reveal that a specific meristematic domain is not utilized in the maize mutant narrow sheath. **Dev Biol** 182: 52–66

Scanlon MJ, Schneeberger RG, Freeling M (1996) The maize mutant narrow sheath fails to establish leaf margin identity in a meristematic domain. **Development** 122: 1683–1691

Schmalfuss K (1930) Untersuchungen über die interkalare Wachstumszone an Glumifloren und dikotylen Bliitenschaften. **Flora** 333–366

Schneeberger RG, Becraft PW, Hake S, Freeling M (1995) Ectopic expression of the *knox* homeobox gene *rough sheath1* alters cell fate in the maize leaf. **Genes Dev** 9: 2292–2304

Schoof H, Lenhard M, Haecker A, Mayer KFX, Jürgens G, Laux T (2000) The stem cell population of *Arabidopsis* shoot meristems is maintained by a regulatory loop between the *CLAVATA* and *WUSCHEL* genes. **Cell** 100: 635–644

Schuster C, Gaillochet C, Lohmann JU (2015) *Arabidopsis* *HECATE* genes function in phytohormone control during gynoecium development. **Development** 142: 3343–3350

Serrano-Mislata A, Bencivenga S, Bush M, Schiessl K, Boden S, Sablowski R (2017) *DELLA* genes restrict inflorescence meristem function independently of plant height. **Nat Plants** 3: 749–754

Serrano-Mislata A, Sablowski R (2018) The pillars of land plants: New insights into stem development. **Curr Opin Plant Biol** 45: 11–17

Sharman BC (1942) Developmental anatomy of the shoot of *Zea mays* L. **Ann Bot** 6: 245–282

Sharman BC, Hitch PA (1967) Initiation of procambial strands in leaf primordia of bread wheat, *Triticum aestivum* L. **Ann Bot** 31: 229–243

Shimizu R, Ji J, Kelsey E, Ohtsu K, Schnable PS, Scanlon MJ (2009) Tissue specificity and evolution of meristematic WOX3 function. **Plant Physiol** 149: 841–850

Shinohara N, Taylor C, Leyser O (2013) Strigolactone can promote or inhibit shoot branching by triggering rapid depletion of the auxin efflux protein PIN1 from the plasma membrane. **PLoS Biol** 11: e1001474

Smith HMS, Hake S (2003) The interaction of two homeobox genes , *BREVIPEDICELLUS* and *PENNYWISE* , regulates internode patterning in the *Arabidopsis* inflorescence. **Plant Cell** 15: 1717-1727

Smith LG, Hake S, Sylvester AW (1996) The tangled-1 mutation alters cell division orientations throughout maize leaf development without altering leaf shape. **Development** 122: 481–489

Snipes SA, Rodriguez K, DeVries AE, Miyawaki KN, Perales M, Xie M, Reddy GV (2018) Cytokinin stabilizes WUSCHEL by acting on the protein domains required for nuclear enrichment and transcription. **PLoS Genet** 14: e1007351

Somssich M, Je B II, Simon R, Jackson D (2016) CLAVATA-WUSCHEL signaling in the shoot meristem. **Development** 143: 3238–3248

Strabala TJ, O'Donnell PJ, Smit AM, Ampomah-Dwamena, C. Martin EJ, Netzler N, Nieuwenhuizen NJ, Quinn BD, Foote HC, Hudson KR (2006) Gain-of-function phenotypes of many *CLAVATA3/ESR* genes, including four new family members, correlate with tandem variations in the conserved *CLAVATA3/ESR* domain. **Plant Physiol** 140: 1331–1344

Suzaki T, Sato M, Ashikari M, Miyoshi M, Nagato Y, Hirano H (2004) The gene *FLORAL ORGAN NUMBER1* regulates floral meristem size in rice and encodes a leucine-rich repeat receptor kinase orthologous to *Arabidopsis* CLAVATA1. **Development** 131: 5649–5657

Suzaki T, Toriba T, Fujimoto M, Tsutsumi N, Kitano H, Hirano HY (2006) Conservation and diversification of meristem maintenance mechanism in *Oryza sativa*: Function of the *FLORAL ORGAN NUMBER2* gene. **Plant Cell Physiol** 47: 1591–1602

Symons GM, Reid JB (2004) Brassinosteroids do not undergo long-distance transport in pea. Implications for the regulation of endogenous brassinosteroid levels. **Plant Physiol** 135: 2196–2206

Takeda T, Suwa Y, Suzuki M, Kitano H, Ueguchi-Tanaka M, Ashikari M, Matsuoka M, Ueguchi C (2003) The *OsTB1* gene negatively regulates lateral branching in rice. **Plant J** 33: 513–520

Tsuda K, Abraham-Juarez M-J, Maeno A, Dong Z, Aromdee D, Meeley R, Shiroishi T, Nonomura K, Hake S (2017) KNOTTED1 cofactors, BLH12 and BLH14, regulate internode patterning and vein anastomosis in maize. **Plant Cell** 29: 1105–1118

Tsuda K, Ito Y, Sato Y, Kurata N (2011) Positive autoregulation of a *KNOX* gene is essential for shoot apical meristem maintenance in rice. **Plant Cell** 23: 4368–4381

Ullah H, Chen J-G, Young J, Im K-H, Sussman M, Jones AM (2001) Modulation of cell proliferation by heterotrimeric G protein in *Arabidopsis*. **Science** 292: 2066–2069

Ung N, Lal S, Smith HMS (2011) The role of PENNYWISE and POUND-FOOLISH in the maintenance of the shoot apical meristem in *Arabidopsis*. **Plant Physiol** 156: 605–614

Urano D, Miura K, Wu Q, Iwasaki Y, Jackson D, Jones AM (2016) Plant morphology of Heterotrimeric G protein mutants. **Plant Cell Physiol** 57: 437–445

Vaughan JG (1955) The morphology and growth of the vegetative and reproductive apices of *Arabidopsis thaliana* (L.) Heynh., *Capsella bursa-pastoris* (L.) Medic. and *Anagallis arvensis* L. **J Linn Soc London Bot** 55: 279–301

Venglat SP, Dumonceaux T, Rozwadowski K, Parnell L, Babic V, Keller W, Martienssen R, Selvaraj G, Datla R (2002) The homeobox gene *BREVIPEDICELLUS* is a key regulator of inflorescence architecture in *Arabidopsis*. **Proc Natl Acad Sci USA** 99: 4730–4735

Vollbrecht E, Veit B, Sinha N, Hake S (1991) The developmental gene *Knotted-1* is a member of a maize homeobox gene family. **Nature** 350: 241–243

Wang W, Li G, Zhao J, Chu H, Lin W, Zhang D, Wang Z, Liang W (2014) DWARF TILLER1, a WUSCHEL-related homeobox transcription factor, is required for tiller growth in rice. **PLoS Genet** 10: e1004154

Webb SE, Appleford NEJ, Gaskin P, Lenton JR (1998) Gibberellins in internodes and ears of wheat containing different dwarfing alleles. **Phytochemistry** 47: 671–677

Wendt T, Holme I, Dockter C, Preuss A, Thomas W, Druka A, Waugh R, Hansson M, Braumann I (2016) HvDep1 Is a Positive Regulator of Culm Elongation and Grain Size in Barley and Impacts Yield in an Environment-Dependent Manner. **PLoS ONE** 11: e0168924

Williams L, Grigg SP, Xie M, Christensen S, Fletcher JC (2005) Regulation of *Arabidopsis* shoot apical meristem and lateral organ formation by microRNA miR166g and its AtHD-ZIP target genes. **Development** 132: 3657–3668

Wolbang CM, Chandler PM, Smith JJ, Ross JJ (2004) Auxin from the developing inflorescence is required for the biosynthesis of active gibberellins in barley stems. **Plant Physiol** 134: 769–777

Wolbang CM, Ross JJ (2001) Auxin promotes gibberellin biosynthesis in decapitated tobacco plants. **Planta** 214: 153–157

Xu C, Liberatore KL, MacAlister CA, Huang Z, Chu YH, Jiang K, Brooks C, Ogawa-Ohnishi M, Xiong G, Pauly M, Van Eck J, Matsubayashi Y, van der Knaap E, Lippman ZB (2015) A cascade of arabinosyltransferases controls shoot meristem size in tomato. **Nat Genet** 47: 784–792

Xu H, Zhao M, Zhang Q, Xu Z, Xu Q (2016) The DENSE AND ERECT

PANICLE 1 (DEP1) gene offering the potential in the breeding of high-yielding rice. **Breed Sci** 66: 659–667

Xu L, Wang C, Cao W, Zhou S, Wu T (2018) CLAVATA1-type receptor-like kinase CsCLAVATA1 is a putative candidate gene for dwarf mutation in cucumber. **Mol Genet Genomics**. 293: 1393-1405

Yadav RK, Girke T, Pasala S, Xie M, Reddy GV (2009) Gene expression map of the Arabidopsis shoot apical meristem stem cell niche. **Proc Natl Acad Sci USA** 106: 4941–4946

Yadav RK, Perales M, Gruel J, Ohno C, Heisler M, Girke T, Jönsson H, Reddy GV (2013) Plant stem cell maintenance involves direct transcriptional repression of differentiation program. **Mol Syst Biol** 9: 654

Yadav RK, Reddy GV (2011) WUSCHEL-mediated cellular feedback network imparts robustness to stem cell homeostasis. **Plant Signal Behav** 6: 544–546

Yadav RK, Tavakkoli M, Xie M, Girke T, Reddy G V. (2014) A high-resolution gene expression map of the *Arabidopsis* shoot meristem stem cell niche. **Development** 141: 2735–2744

Yanai O, Shani E, Dolezal K, Tarkowski P, Sablowski R, Sandberg G, Samach A, Ori N (2005) *Arabidopsis* KNOX1 proteins activate cytokinin biosynthesis. **Curr Biol** 15: 1566–1571

Yokoyama R, Takahashi T, Kato A, Torii KU, Komeda Y (1998) The Arabidopsis ERECTA gene is expressed in the shoot apical meristem and organ primordia. **Plant J** 15: 301–310

Yoon J, Cho L-H, Antt HW, Koh H-J, An G (2017) KNOX protein OSH15 induces grain shattering by repressing lignin biosynthesis genes. **Plant Physiol** 174: 312 LP-325

Yoshikawa T, Tanaka S-Y, Masumoto Y, Nobori N, Ishii H, Hibara K-I, Itoh J-I, Tanisaka T, Taketa S (2016) Barley *NARROW LEAFED DWARF1* encoding a WUSCHEL-RELATED HOMEBOX 3 (WOX3) regulates the marginal development of lateral organs. **Breed Sci** 66: 416–424

Zhang H, Wu K, Wang Y, Peng Y, Hu F, Wen L, Han B, Qian Q, Teng S (2012) A WUSCHEL-like homeobox gene, *OsWOX3B* responses to NUDA/GL-1 locus in rice. **Rice** 5: 30

Zhou W, Malabanan PB, Abrigo E (2015) OsHox4 regulates GA signaling by interacting with DELLA-like genes and GA oxidase genes in rice. **Euphytica** 201: 97–107

Zhu Y, Nomura T, Xu Y, Zhang Y, Peng Y, Mao B, Hanada A, Zhou H, Wang R, Li P, et al (2006) *ELONGATED UPPERMOST INTERNODE* encodes a cytochrome P450 monooxygenase that epoxidizes gibberellins in a novel deactivation reaction in rice. **Plant Cell** 18: 442–456

Figure 1. Proliferative patterns associated with axial elongation

(A) Arabidopsis vegetative shoot apical meristem and rosette. Proliferation in the apical central zone displaces cells into the neighbouring peripheral zones. Leaf primordia form off the peripheral zone. Underlying rib meristem produces files of transversely dividing cells which together with subapical peripheral cells, displace downwards to form the stem. Little internode elongation between nodes leads to a rosette habit. (B) Vegetative shoot apical meristem and compressed seedling modelled from maize. Similar to (A), the meristem has central, proliferative and rib zones and produces leaf primordia off its peripheral flank; however, leaf primordia remain attached to the meristem base by a disc of insertion (doi). Clonal analyses shows that the upper doi forms the leaf blade while the lower doi will develop into the node, internode and axillary bud. Little internode elongation between nodes leads to a compressed stacked internodes. (C) Bolting Arabidopsis inflorescence and stem. The rib meristem increases in size and activity to fuel acropetal expansion of internodes between secondary axillary buds (paraclade) and later to separate flower-bearing pedicels. (D) Grass culm elongation following flowering modelled from barley. Internode elongation occurs in an acropetal pattern up the culm. Division in intercalary meristems displaces cells upwards to expand and mature. The intercalary meristem eventually arrests and all internode cells are expanded. Gray shading indicates the rib zone region of stem morphogenesis. Pale pink shading shows floral meristems produced off the flanks of the inflorescence meristem. Peach colour shows active intercalary meristem fueling grass internode growth. Pale green denotes leaves and leaf primordia. CZ, central zone; rm, rib meristem; im, intercalary meristem; doi, disc of insertion

Figure 2. Pathways and expression patterns associated with subapical proliferation and intercalary meristems

(A) An Arabidopsis apical meristem. Two overlapping pathways regulate meristem function at the shoot apex. The negative feedback between *WUSCHEL* (*WUS*) expression in the rib meristem and *CLAVATA3* (*CLV3*) peptide production in the overlying cells acts to maintain meristem integrity, size and proliferation rate, thus controlling the cells available for stem morphogenesis. Disruptions in this loop cause stem fasciation, associated with altered cell division orientations and dwarfism. *WUS* expression is reinforced by localised accumulation of cytokinin. The Class-I KNOX genes (*SHOOT MERISTEMLESS* (*STM*), *BREVIPEDICELLUS* (*BP*) and *KNOTTED-like HOMEBOX2* (*KNOX2*)) promote indeterminacy in the meristem, downregulate GA biosynthesis and are themselves downregulated in emerging lateral primordia. *KNOX2* and *BP* are expressed more strongly at the meristem boundaries and in the rib zone. (B) A grass apical meristem modelled on maize. The basal boundary of *ZmWUS1* expression in the central meristem is limited by FEA3 perception of the FCP1 (FON2-LIKE CLE PROTEIN1), a *CLAVATA3* /ENDOSPERM SURROUNDING REGION (CLE) peptide produced by the leaf primordium. Expression of *NARROW SHEATH* (*NS*) in the marginal domain of the leaf primordium and underlying disc of insertion is important to recruit founder cells for leaf margins and internodes. *ZmWUS1* reappears in the lower disc expanding towards the presumptive

axillary bud. *NS* genes in rice repress GA biosynthesis, and may do the same in maize. *Class-I KNOX* genes are expressed in the apical meristem, with *KNOTTED1* (*Kn1*) has the broadest expression through the meristem while *Rough sheath1* (*Rs1*) and *GNARLY1* (*Gn1*) are expressed more predominantly in the disc of insertion. All three remain expressed in the presumptive internode region and vasculature. **(C)** Maize early intercalary meristem. *KN1* remains expressed in the intercalary meristem. *Rs1* and *Gn1* expression has not been examined for intercalary meristem tissue. It is also unknown whether there is a CLE peptide – CLV receptor which regulates intercalary meristem size. Expression of the *NS WOX3b* sister group is observed in maize and rice.

Figure 3. Control of boundary gene expression in the rib zone and intercalary meristem and internode

(A) Arabidopsis rib zone shortly after the reproductive transition. Expression of *PENNYWISE* (*PNY*) in the rib meristem promotes transverse cell divisions and limits expression of boundary genes to the rib zone periphery. *PNY* physically interacts with *BREVIPEDICELLUS* (*BP*) where both act to exclude boundary gene expression from the developing internode and repress lignin biosynthetic gene expression. Gray shading indicates the rib zone region. Pale pink shading shows floral meristems. **(B)** *BEL-LIKE HOMEODOMAIN12* (*BLH12*)/*BLH14* are expressed in the intercalary meristem of maize where they are essential to promote meristem maintenance, as well as the internode vasculature (purple stripes) where they prevent premature differentiation and anastomoses. *BLH12/14* directly interacts with *KNOTTED1* (*KN1*) which is also expressed in the intercalary meristem and vasculature, although the relevance of this interaction to *BLH12/14* function is unknown. Peach colour shows active intercalary meristem.




Figure 1




Figure 2

